2019 STAAR Algebra I Rationales

Item #		Rationale
1	Option D is correct	To determine the slope (steepness) of a straight line when graphed on a coordinate grid; $m = \frac{y_2 - y_1}{x_2 - x_1}$); of the line represented by the equation $y = 12x - 19$, the student could have recalled the slope-intercept form of a linear equation, $y = mx + b$, where m represents the slope and b represents the y-intercept (value where the line crosses the y-axis). Therefore, the slope of the graph is represented by the value of 12. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely identified the value of b in the slope-intercept form of a linear equation, $y = mx + b$, as the slope of the line represented by the graph. The student needs to focus on understanding the key parts of the slope-intercept form of a linear equation.
	Option B is incorrect	The student likely identified the slope of the line represented by the graph as the quotient (answer to a division problem) of the coefficient (a number used to multiply a variable) of the x term divided by the value of the y-intercept. The student needs to focus on understanding the key parts of the slope-intercept form of a linear equation.
	Option C is incorrect	The student likely identified the slope of the line represented by the graph as the value of the x-intercept (value where a line crosses the x-axis). To calculate the x-intercept, the student set y equal to 0, added 19 to both sides of the equation, and divided both sides of the equation by 12 to solve for x. The student needs to focus on understanding the key parts of the slope-intercept form of a linear equation.

Item #		Rationale
3	Option D is correct	To determine what value of x makes the given equation true, the student could have solved the equation for x. To solve the equation, the student could have first distributed (multiplied) the 1.25 in front of the parentheses by the values inside the parentheses. This step results in the equation $1.25(4x) + 1.25(10) = 7.5$, which is $5x + 12.5 = 7.5$. The student could have then added -12.5 to both sides of the equation, resulting in $5x = -20$. To solve for x, the student could have divided by 5 on both sides of the equation, resulting in $x = -4$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely did not distribute the 1.25 on the left side of the equation to the second term inside the parentheses, 10, resulting in the equation $1.25(4x) + 10 = 7.5$, which is $5x + 10 = 7.5$. The student likely added -10 to both sides of the equation, resulting in $5x = -17.5$. To solve for x, the student likely divided by 5 on both sides of the equation, resulting in $x = -3.5$. The student needs to focus on applying the distributive property correctly when solving equations.
	Option B is incorrect	The student likely distributed correctly but eliminated 12.5 from the left side of the equation by subtracting 12.5 on the right side of the equation, resulting in the equation $5x = -45$. To solve for x, the student likely divided by 5 on both sides of the equation, resulting in $x = -41$. The student needs to focus on isolating the variable correctly when solving equations.
	Option C is incorrect	The student likely did not distribute the 1.25 on the left side of the equation to the second term inside the parentheses, 10, resulting in the equation $1.25(4x) + 10 = 7.5$, which is $5x + 10 = 7.5$. The student likely subtracted 10 from both sides of the equation, resulting in $5x = -2.5$. To solve for x, the student likely divided by 5 on both sides of the equation, resulting in $x = -0.5$. The student needs to focus on applying the distributive property and isolating the variable correctly when solving equations.

Item#		

Item#	Rationale	
6	Option F is correct.	To determine the equation of a line in slope-intercept form (y = mx + b)
		4
		4
		4
		4

Item #		Rationale
7	Option C is correct	To determine which graph best represents the equation $45y = 46x + 15$, the student could have first isolated the y on one side of the equation by dividing both sides of the equation by 45, resulting in $y = \frac{6}{5}x + \frac{1}{3}$. Next, the student could have verified that the line will cross the y-axis (vertical) at $(0, \frac{1}{3})$ and

Item #	Rationale	
9	Option A is correct	To determine which function is represented by the graph, the student could have determined that the graph can be represented by an exponential function, $g(x) = ab^x$, where a is the y-intercept (value where the graph crosses the y-axis), b is the decay factor (constant rate that the value is decreased by over a period of time), and x is the variable (symbol used to represent an unknown number). To write the exponential function, the student could have identified 6 as the y-intercept. Next, the student could have determined the decay factor as $\frac{1}{3}$ by dividing each y-coordinate by the previous y-coordinate. The ordered pairs $(4, 18)$, $(6, 6)$, and $(12, 2)$; $6 \div 18 = \frac{1}{3}$ and $2 \div 6 = \frac{1}{3}$. The
		4
		4

Item #	Rationale	
10	Option J is correct	To determine the equation of a line perpendicular (a line meeting another line at a right angle, 90°) to another line, the student could have recalled that the slope (steepness of a straight line) when

Item #	Rationale	
	Option H is incorrect	The student likely identified the slope of the line perpendicular to the given line correctly as $\frac{9}{4}$ but used $4\frac{9}{4}$ to determine the y-intercept of the line perpendicular to the given line. The student needs to focus on understanding how to determine the slope of a line perpendicular to a given line.

Item #	Rationale	
11	Option C is correct	To determine the domain (all possible x-values) of the function (a relationship where each input has a single output), the student could have identified all the x-values for which the graph has a y-value. The graph goes from 0 on the left to 50 on the right, making the domain all real numbers greater than or equal to 0 and less than or equal to 50. To determine the range (all possible y-values), the student could have identified all the y-values for which the graph has an x-value. The graph goes from 0 on the lowest point to 100 on the highest point, making the range all real numbers greater than or equal to 0 and less than or equal to 100.
	Option A is incorrect	The student likely reversed the relationship between domain and range. The student needs to focus on understanding how to identify the domain and range from the graph of a function.
	Option B is incorrect	The student likely identified domain as the slope (steepness of a straight line when graphed on a coordinate grid; $m = \frac{y_2 - y_1}{x_2 - x_1}$) and range as the y-intercept (value where a line crosses the y-axis) of the line representing the function. The student needs to focus on understanding how to identify the domain and range from the graph of a function.
	Option D is incorrect	The student likely identified domain as the y-intercept (value where a line crosses the y-axis) and range as the slope (steepness of a straight line when graphed on a coordinate grid; $m = \frac{y_2 - y_1}{x_2 - x_1}$) of the line representing the function. The student needs to focus on understanding how to identify the domain and range from the graph of a function.

Item#	Rationale	
13	Option B is correct	To determine the equivalent expression, the student could have divided 45 by 15 and used the quotient of powers property $\frac{a^m}{a^n} = a^{(m-n)}$, resulting in $3m^{(6-4)}p^{(2-8)}v^{(12-(-4))} = m^{(6-(-2))}p^{(2-8)}v^{(12+4)} = 3m^{8}p^{-6}v^{16}$. Then the student could have used the negative exponent property $a^{-n} = \frac{1}{a^n}$, resulting in $3\frac{1}{m^4}\frac{1}{p^6}v^{16} = \frac{3v^{16}}{m^4p^6}$. The rationale for the correct answer is that could wey2 (the)64o2 (the)6sol e.EMC /TD <</MHID 40 >>BC 0 0 0 -1 Tm7 Td4<00B7>[(Ope)]T(properA

2019 STAAR Algebra I Rationales

Item#	Rationale	
15	Option B is correct	To determine the inequality representing all possible combinations of x, the number of sausage links, and y, the number of briskets cooked on the grill, the student could have first identified that each sausage link weighed 2 pounds and each brisket weighed 6 pounds, represented by the expressions $2x$ and $6y$. So the total number of pounds of sausage links and briskets cooked on the grill could be represented by the expression $2x + 6y$. Then the student could have realized the phrase "no more than" can be represented by the inequality symbol $\leq$, so the phrase "no more than 120 lb" can be represented by ≤ 120 lb. Therefore all possible combinations of x and y could be represented by the inequality $2x + 6y \leq 120$. The rationale for the correct answer is one efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student reversed the coefficients (a number used to multiply a variable) representing the weights of each sausage link and each brisket and identified the phrase "no more than" as being represented by the inequality symbol $<$, resulting in $6x + 2y < 120$. The student needs to focus on understanding how to write a linear inequality given a real-world situation.
	Option C is incorrect	The student reversed the coefficients (a number used to multiply a variable) representing the weights of each sausage link and each brisket and identified the phrase "no more than" as being represented by the inequality symbol $>$, resulting in $6x + 2y > 120$. The student needs to focus on understanding how to write a linear inequality given a real-world situation.
	Option D is incorrect	The student identified the coefficients (a number used to multiply a variable) representing the weights of each sausage link and each brisket correctly but identified the phrase "no more than" as being represented by the inequality symbol is4d78ug0

2019 STAAR Algebra I Rationales

Item#	Rationale	
17	Option D is correct	To determine which graph represents the solution set of $y > 3x + 4$, the student could have determined that the $>$ symbol means "greater than," which can be represented by a dashed line (to indicate no points on the line are part of the solution set) and a shaded area of the graph. The dashed line should cross the y-axis (vertical axis) at 4.

Item#		Rationale
	Option C is incorrect	The student correctly identified a line with the correct slope (3) and y-intercept (44). The student identified the inequality symbol > as meaning less than, which could be represented by a dashed line (to indicate no points on the line are part of the solution set) and the shaded part of the graph that does not include the origin (0, 0), interpreting 0 > 44 as not being a true statement. The student needs to focus on understanding how inequality symbols affect the graph of the solution set of a linear inequality.

Item #	Rationale	
18	Option G is correct	To determine the best interpretation of one of the values in the function $b(x) = 850(1.025)^x$, the student could have recognized that an exponential function written in the form $b(x) = ac^x$, a represents the initial value, c is 1 plus the rate that the value is increased (when $c > 1$) or 1 minus the factor that the value is decreased (when $c < 1$), and x is the variable (symbol used to represent an unknown number). In this situation, the variable x represents the number of years. In $b(x) = 850(1.025)^x$, 850 is the initial value, and because $1.025 > 1$, 0.025 represents an increase rate of 0.025 each year, or 2.5% each year. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect	The student likely identified the rate in the situation as $1 - 0.025 = 0.975$, which represents a decrease rate of 97.5% each year. The student needs to focus on interpreting the meaning of the values of a and c of exponential functions in the form $b(x) = ac^x$.
	Option H is incorrect	The student likely identified the value of c as the initial balance and misidentified the decimal as a comma (1,025). The student needs to focus on interpreting the meaning of the values of a and c of exponential functions in the form $b(x) = ac^x$.
	Option J is incorrect	The student likely identified the phrase "at the end of one year" as representing the initial value (850) of the exponential function. The student needs to focus on interpreting the meaning of the values of a and c of exponential functions in the form $b(x) = ac^x$.

		4 4

Item #		Rationale
21	Option A is correct	To determine which graph best represents the equation $y = 4x^2 + 6x - 41$, the student could have first identified the equation as being in the standard form of a quadratic equation ($y = ax^2 + bx + c$); therefore $a = 4$, $b = 6$, and $c = -41$. Then the student could have solved for the axis of symmetry $\left(x = \frac{-b}{2a}\right)$ of the graph, resulting in $x = \frac{-6}{2(4)} = \frac{-6}{8} = -\frac{3}{4}$. Then the student could have solved for the y-value of the vertex (highest or lowest point of the curve) by substituting $x = -\frac{3}{4}$ into the equation, resulting in $y = 4(-\frac{3}{4})^2 + 6(-\frac{3}{4}) - 41 = -\frac{89}{4}$. Then the student could have solved for the y-intercept (value where graph crosses the y-axis) by substituting $x = 0$ into the equation, resulting in $y = 4(0)^2 + 6(0) - 41 = -41$. Lastly, the student could have identified the parabola (U-shaped graph) that has an axis of symmetry at $x = -\frac{3}{4}$ and a vertex at $(-\frac{3}{4}, -\frac{89}{4})$ and that crosses the y-axis (vertical) at -41. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option B is incorrect	The student likely graphed a parabola with the correct y-intercept but with an axis of symmetry at $x = -\frac{4}{3}$ and a vertex at $(-\frac{4}{3}, -\frac{410}{3})$. The student needs to focus on understanding how to graph a quadratic function on the coordinate plane and identifying key features of the graph.
	Option C is incorrect	The student likely graphed a parabola that has an axis of symmetry at $x = -\frac{4}{3}$ and a vertex at $(-\frac{4}{3}, -\frac{8}{3})$ and that crosses the y-axis at -41. The student needs to focus on understanding how to graph a quadratic function on the coordinate plane and identifying key features of the graph.
	Option D is incorrect	The student likely graphed a parabola that has the correct axis of symmetry, but with a vertex at $(3, 10)$, and that crosses the y-axis at -41. The student needs to focus on understanding how to graph a quadratic function on the coordinate plane and identifying key features of the graph.

Item #		Rationale
22	Option H is correct	To determine the best prediction of the number of clicks on the advertisement if 1,500 people visited the website, the student could have first used a graphing calculator to generate the function using linear regression (a method of determining the line of best fit). The function that best models the data is $y = 0.0519x + 5.5393$. Next, the student could have substituted 1,500 for x in the function and solved for y, resulting in

Item#	Rationale
	Option J is incorrect. The student likely multiplied each value of the ordered pair (307, 21) by 5, since the product (answer) of 307 and 5 is 1,535, which is closest to 1,500. The product of 21 and 5 is 105. The student needs to focus on understanding how to make predictions for real-world problems represented by linear functions.

Item #		Rationale
24	Option H is correct	To determine the equivalent expression, the student could have applied the power of a power property $\left((a^m)^n = a^{mn}\right)$, resulting in $x^{1 \cdot 2} y^{4 \cdot 6 \cdot 2} = x^2 y^{4 \cdot 12}$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect	The student squared (multiplied a number by itself) the exponents (numbers raised to a power) 1 and 4(6) inside the parentheses, resulting in $x^{(1)^2} y^{4(6)^2} = 1x y^{436} = xy^{436}$. The student needs to focus on understanding how to use the properties of exponents to simplify expressions with powers raised to a power.
	Option G is incorrect	The student squared (multiplied a number by itself) the exponents (numbers raised to a power) 1 and 46 inside the parentheses, resulting in $x^{(1)^2} y^{(6)4^2} = x^{136} = xy^{36}$. The student needs to focus on understanding how to use the properties of exponents to simplify expressions with powers raised to a power.
	Option J is incorrect	The student did not include the negative sign with 6 when multiplying the exponents (numbers raised to a power) by the variable (symbol used to represent an unknown number) y, resulting in $x^{1 \cdot 2} y^{6 \cdot 2} = x^2 y^{12}$. The student needs to focus on understanding how to use the properties of exponents to simplify expressions with powers raised to a power.

Item #	Rationale	
25	Option A is correct	To determine the number of courses completed worth 3 credits, the student could have set up and solved a system of linear equations (two or more linear equations containing the same set of variables (symbols used to represent unknown numbers)). Using the variables t as the number of courses completed worth 3 credits and f as the number of courses completed worth 4 credits, the equation for the total number of courses completed is $t + f = 18$, and the equation relating to the total number of credits earned is $3t + 4f = 59$. Next, the student could have subtracted t from both sides of the first equation to solve for f, resulting in $f = 18 - t$. Using the substitution method, the student could have substituted $f = 18 - t$ into the second equation, resulting in $3t + 4(18 - t) = 59$. The student could have then distributed (multiplied) the 4 to the terms inside the parentheses, resulting in $3t + 72 - 4t = 59$. The student could have then combined like terms (terms that contain the same variables raised to the same powers), resulting in $72 - t = 59$. The student could have then subtracted 72 from both sides of the equation, resulting in $-t = -13$. Lastly, the student could have divided both sides of the equation by -1, resulting in $t = 13$. Therefore, the number of courses completed worth 3 credits is 13. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option B is incorrect	The student likely set up and solved the system of linear equations for the number of courses completed worth 4 credits instead of the number of courses completed worth 3 credits. The student needs to focus on understanding how to interpret and solve a system of equations given verbal descriptions.
	Option C is incorrect	The student likely set up and solved the system of linear equations for the number of courses completed worth 4 credits and then calculated the total number of credits earned from the courses worth 4 credits ($4f = 4 \cdot 5 = 20$). The student needs to focus on understanding how to interpret and solve a system of equations given verbal descriptions.
	Option D is incorrect	The student likely set up and correctly solved the system of linear equations for the number of courses completed worth 3 credits but then calculated the total number of credits earned from the courses worth 3 credits ($3t = 3 \cdot 13 = 39$). The student needs to focus on understanding how to interpret the verbal descriptions as a system of equations.

Item #	Rationale	
26	Option J is correct	To determine the zero (input value, x) that produces an output value, y, of 0 for a linear function f, the student could have plotted the point (1, 49) on the coordinate grid and drawn a line that goes through that point and has a slope (steepness of a line when graphed on a coordinate grid; $m = \frac{y_2 - y_1}{x_2 - x_1}$) of 43. Then the student could have identified the x-value where the line crosses the x-axis (horizontal), which is 42. Therefore, the zero of f is 42. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect	The student likely drew a line with the correct slope but through a point plotted at (41, 9) instead of (1, 49), resulting in the line crossing the x-axis at 2. The student needs to focus on understanding how to plot points on a coordinate grid and identifying key features of linear functions.
	Option G is incorrect	The student likely drew a line with the correct slope but through a point plotted at (1, 9) instead of (1, 49), resulting in the line crossing the x-axis at 4. The student needs to focus on understanding how to plot points on a coordinate grid and identifying key features of linear functions.
	Option H is incorrect	The student likely interpreted the zero of the function as representing the y-intercept (value where a line crosses the y-axis) of the line. The student needs to focus on understanding how to identify key features of linear functions.
27	29 and any equivalent values are correct	The student could have realized $h(x) = 29(5.2)^x$ represents an exponential function, $h(x) = ab^x$, where a is the y-intercept (value where a graph crosses the y-axis), b is the growth factor, and x is the variable (symbol used to represent an unknown number). Therefore, the value of the y-intercept of the graph of $h(x) = 29(5.2)^x$ is 29. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.

Item #		Rationale
28	Option H is correct	To determine which function is best represented by the graph, the student could have identified the solutions (x-values when y is equal to 0) of the function as u and v and used the solutions to construct and simplify the equation of a quadratic function using $h(x) = (x - u)(x - v)$. The solutions can be identified by where the parabola (U-shaped graph) crosses the x-axis (at $x = 0$ and $x = 6$). Letting $u = 0$ and $v = 6$, the student could have substituted those values into the equation $h(x) = (x - u)(x - v)$, resulting in $h(x) = (x - 0)(x - 6)$. The student could have then multiplied the expressions $(x - 0)$ and $(x - 6)$, resulting in $h(x) = x^2 - 6x - 0x + 0$. Lastly, the student could have combined like terms (terms that contain the same variables raised to the same powers), resulting in $h(x) = x^2 - 6x$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect	The student likely identified the solutions of the function as the x- and y-coordinates of the vertex but identified the ordered pair of the vertex (high or low point of the curve) as $(-3, -49)$ instead of $(3, -49)$. The student likely substituted those values for b and c in the standard form of a quadratic equation $\left(h(x) = ax^2 + bx + c\right)$, resulting in $h(x) = x^2 - 3x - 9$. The student needs to focus on understanding how to identify the solutions of a quadratic function and write the equation of the function using those solutions.
	Option G is incorrect	The student likely identified the solutions of the function as the x- and y-coordinates of the vertex (high or low point of the curve), $(3, -49)$, and substituted those values for b and c in the standard form of a quadratic equation $\left(h(x) = ax^2 + bx + c\right)$, resulting in $h(x) = x^2 + 3x - 9$. The student needs to focus on understanding how to identify the solutions of a quadratic function and write the equation of the function using those solutions.
	Option J is incorrect	The student identified the correct solutions of the function but substituted the solutions into the equation $h(x) = (x + u)(x + v)$, resulting in $h(x) = (x + 0)(x + 6)$. The student likely multiplied the expressions $(x + 0)$ and $(x + 6)$, resulting in $h(x) = x^2 + 6x + 0x + 0$. Lastly, the student likely combined like terms, resulting in $h(x) = x^2 + 6x$. The student needs to focus on understanding how to write the equation of a quadratic function using the solutions of the function.

Item #	Rationale	
29	Option C is correct	To determine the equivalent expression, the student could have found the factors (numbers or expressions that can be multiplied together to get another number or expression) of $24x^2 - 22x + 5$. To find the factors, the student could have determined that $24x^2$ is equal to $12x \cdot 2x$ and written $12x$ and $2x$ as the first terms in each factor. The student could have then determined that the second terms of the factors are -5 and -1 because their product (answer when multiplying) is 5 (last term in the expression given), and $(12x \cdot (-1)) + (-10 \cdot 2x) = -12x - 10x = -22x$ (number in the middle term in the expression given). The student could have then written the two factors as $(12x - 5)(2x - 1)$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely verified that $12x \cdot 2x = 24x^2$ and $5 \cdot 1 = 5$ but did not verify the middle term of the expression: $(12x \cdot 1) + (5 \cdot 2x) = 12x + 10x = 22x$. The student needs to focus on understanding how to factor trinomials in the form $ax^2 + bx + c$.
	Option B is incorrect	The student likely verified that $8x \cdot 3x = 24x^2$ and $-5 \cdot (-1) = 5$ but did not verify the middle term of the expression: $(8x \cdot (-1)) + (-5 \cdot 3x) = -8x - 15x = -23x$. The student needs to focus on understanding how to factor trinomials in the form $ax^2 + bx + c$.
	Option D is incorrect	The student likely verified that $8x \cdot 3x = 24x^2$ and $5 \cdot 1 = 5$ but did not verify the middle term of the expression: $(8x \cdot 1) + (5 \cdot 3x) = 8x + 15x = 23x$. The student needs to focus on understanding how to factor trinomials in the form $ax^2 + bx + c$.

Item #	Rationale	
	Option H is incorrect	The student likely calculated the slope of each line correctly but identified the value of each b as the x-intercept (value where a line crosses the x-axis) ($b_1 = 44$ and $b_2 = 42$), resulting in the equations $y = x 44$ and $y = 42x 42$. The student needs to focus on understanding how to write a system of two linear equations in slope-intercept form when given a graph.

Item#	Rationale	
31	Option B is correct	To determine the function, the student could have determined that the relationship can be represented by an exponential function, $f(x) = ab^x$, where a is the initial value, b is the common factor, and x represented B. Hmhe992 ((symbJ /TT1 1 Tf 4.136 0 Td (a)Tj /T1_0 1 Tf 0.951 0 TdTf 0.943 0 f 5.684 0 Td (x)Tj /T

Item #	Rationale	
	Option D is incorrect	The student likely identified the initial value correctly but determined the common factor by dividing each $f(x)$-value by the next $f(x)$-value $\left(\frac{12.5}{15} = \frac{5}{6}, \frac{15}{18} = \frac{5}{6}, \frac{18}{21.6} = \frac{5}{6}, \frac{21.6}{25.92} = \frac{5}{6}\right)$, resulting in the exponential function $f(x) = 18\left(\frac{5}{6}\right)^x$. The student needs to focus on understanding how to find the common factor of an exponential function from a given table of values.

Item#		Rationale
33	Option D is correct	To determine the best statement about the transformation of the graph of $f(x) = x^2$ to the graph of $g(x) = f(x) - 4.9$, the student could have recognized that subtracting 4 from $f(x)$
		4
		4
		4

2019 STAAR Algebra I Rationales

Item #	Rationale	
35	Option D is correct	To determine which statement is true about linear function k, the student could have plotted the points $(4, 7)$ and $(1, 8)$ on the coordinate grid and drawn a line passing through the two points, recognizing the x-intercept (value where line crosses the x-axis) of the graph as 4. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely made a sign error when calculating the slope (steepness) of a straight line when graphed on a coordinate grid; $m = \frac{y_2 - y_1}{x_2 - x_1}$ of the line, resulting in $m = \frac{8 - 0}{1 - 7} = \frac{8}{-6} = -\frac{4}{3}$. The student needs to focus on understanding the key features of linear functions graphed on a coordinate grid.
	Option B is incorrect	The student identified a point representing the opposite values of the x- and y-coordinates of the point $(1, 8)$ as being on the graph of linear function k. The student needs to focus on understanding the key features of linear functions graphed on a coordinate grid.
	Option C is incorrect	The student identified the zero (value of x when $y = 0$) as the y-intercept (value where line crosses the y-axis) of linear function k. The student needs to focus on understanding the key features of linear functions graphed on a coordinate grid.

Item #	Rationale	
36	Option F is correct	To determine the equivalent expression, the student could have realized that $21d$ is the common factor (a number or expression that divides another number or expression) of $210d^2$ and $463d$. Because $21 \cdot 10 = 210$ and $d \cdot d = d^2$, $210d^2$ can be expressed as $21(10)d$. Similarly, $463d$ can be expressed as $21d(43)$. The student could have recognized the common factor as $21d$ and subtracted the other two terms to indicate that $210d^2 - 463d$

Item#	Rationale	
37	Option C is correct	To determine the x-value of the solution to the system of equations (two or more equations containing the same set of variables (symbols used to represent unknown numbers)), the student could have used the substitution method to find the value of x. First, the student could have substituted the second equation, $y = 45x + 32$, for y in the first equation, $3x + 45y = 22$, resulting in $3x + 45(54x + 32) = 22$. Then, the student could have distributed (multiplied) 45 to both terms inside the parentheses, resulting in $3x + 2430x + 1440 = 22$. Then, the student could have combined the x-terms, resulting in $2433x + 1440 = 22$. Then, the student could have added 1440 to both sides of the equation, resulting in $2433x = 1418$. Lastly, the student could have divided by 2433 on both sides of the equation, resulting in $x = 0.583$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely made a sign error when subtracting 160 from both sides of the equation, $28x + 160 = 22$, resulting in $28x = 182$. The student needs to focus on understanding how to complete all the steps to calculate the solution to a system of equations.
	Option B is incorrect	The student likely made sign errors and calculated the $y = 4x$ y-value instead of the x-value in the solution to the system of equations,

Item#	Rationale	
38	Option G is correct	To determine which graph best represents the equation, the student could have calculated the values of y for several values of x and determined which graph includes those values. Using the values of 0, 1, and 2 for x yields the points (0, 10), (1, 18.5), and (2, 27.225), which are all on this graph. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect	The student likely made sign errors on the x-values when identifying the points on the coordinate grid, using the points (−1, 8.5) and (−2, 7.225). The student needs to focus on understanding how to identify graphs of exponential equations.
	Option H is incorrect	The student likely added 0.85 to the given equation, identifying the graph of $y = 10(1.85)^x$. The student needs to focus on understanding how to identify graphs of exponential equations.
	Option J is incorrect	The student likely subtracted 0.85 from the given equation from 1, identifying the graph of $y = 10(0.15)^x$. The student needs to focus on understanding how to identify graphs of exponential equations.

Item#	Rationale	
40	Option J is correct	To determine which value of x is a solution to the equation, the student could have found the factors (numbers or expressions that can be multiplied together to get another number or expression) of $5x^2 + 36x + 36$ and solved the factors for x. The student could have determined that $5x^2$ is equal to $5x \cdot x$ and written $5x$ and x as the first terms in the factors. The student could have then determined that the second terms of the factors are 6 and 6 because their product (answer when multiplying) is 36 (last term in the expression), and

2019 STAAR Algebra I Rationales

Item #	Rationale	
41	Option A is correct	To determine the domain (all possible x-values) of the part of the exponential function graphed on the grid, the student could have identified all the values of x for which the graph has a y-value. The graph extends upward indefinitely from -2 on the left, making the domain $x \geq -2$.
	Option B is incorrect	The student likely interpreted domain (Ž) to mean all possible y-values instead of all possible x-values. The student likely identified the graph as extending upward indefinitely from -4.5 on the bottom, making the domain $y \geq -4.5$. The student needs to focus on understanding how to represent the domain from a graph of an exponential function.
	Option C is incorrect	The student likely identified the variable (symbol used to represent an unknown number) representing the domain correctly but interpreted domain (Ž) to mean all possible y-values instead of all possible x-values. The student likely identified the graph as extending upward indefinitely from -4.5 on the bottom, making the domain $x \geq -4.5$. The student needs to focus on understanding how to represent the domain from a graph of an exponential function.
	Option D is incorrect	The student likely determined the domain correctly but identified the variable (symbol used to represent an unknown number) y instead of x as representing the domain, making the domain $y \geq -2$. The student needs to focus on understanding how to represent the domain from a graph of an exponential function.
42	-38 and any equivalent values are correct	To determine the solution, the student could have solved the equation for m. To solve the equation, the student could have first distributed (multiplied) -4 to the terms inside the first set of parentheses and 4 to the terms inside the second set of parentheses, resulting in $4(6m) + (-4)(8) = 4(17) + 4(4m) \Rightarrow 46m - 48 = 68 - 4m$. The student could have then added 48 to both sides of the equation, resulting in $46m = 76 - 4m$. The student could have then added $4m$ to both sides of the equation, resulting in $42m = 76$. To solve for m, the student could have divided both sides of the equation by 42, resulting in $m = \frac{38}{21}$. Therefore, the solution to the equation is $\frac{38}{21}$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.

Item#	Rationale	
43	Option B is correct	To determine which function is equivalent to $y = 3(x + 2)^2 + 7$, the student could have simplified the equation in ch978 0 0/T1_1 1 Tf 0 Tc 1.811 0 Td (+)Tj /T1_0 1 Tf nd26.8 (v) [(equatioform[(equatioof[(equatioa[(equatio Tfd (ation

Item#	Rationale	
44	Option H is correct	To determine which table shows y as a function of x, the student could have recalled that a function represents a relationship where each input, x, has a single output, y. Also, the student could have recalled that a function can have repeated output values as long as the input values are different. A student could have analyzed the ordered pairs (1, 4), (3, 4), (7, 4), and (12, 4) to conclude that each x-value has a single y
	4 4	

		4
		44

2019 STAAR Algebra I Rationales

Item #	Rationale	
47	Option A is correct	To determine which graph best represents the solution set (set of all numbers that satisfy an equation

Item #		Rationale
	Option D is incorrect	The student likely identified the reciprocal (pair of numbers that, when multiplied together, equal 1) of both slopes of the lines and solved $8x = 16$ for the y-intercept of the line representing the first equation and $3x = 6$ for the y-intercept of the line representing the second equation. The line of the first equation has a slope of $\frac{1}{2}$ and a y-intercept of 2, and the line of the second equation has a slope of -45 and a y-intercept of 2, resulting in the equations $y = \frac{1}{2}x + 2$ and $y = -45x + 2$. The student needs to focus on understanding how to set up linear equations and graph a system of two linear equations on a coordinate grid.

Item #	Rationale	
50	Option F is correct	To determine the domain (all possible x-values) and range (all possible y-values) of g(x), the student could have graphed the quadratic function on a coordinate grid and analyzed the parabola (U-shaped graph). The student could have determined that the graph continues to extend downward and outward indefinitely, making the domain all real numbers. As the graph of this function opens downward, it is bound above by y = 61, making the range all values of y less than or equal to 61 (y 61), or g(x) 61. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option G is incorrect	The student identified the range correctly by using the x-value of the vertex (high or low point of the curve) as the upper bound of the domain, x 17. The student needs to focus on understanding how to identify the domain of a quadratic function.
	Option H is incorrect	The student identified the domain correctly by identified the range as all possible x-values with an upper bound represented by the x-value of the vertex (high or low point of the curve), x 17. The student needs to focus on understanding how to identify the range of a quadratic function.
	Option J is incorrect	The student likely used the x-value of the vertex (high or low point of the curve) as the upper bound of the domain, x 17, and the y-value of the vertex as the lower bound of the range, g(x) 61. The student likely then reversed the relationship between domain and range. The student needs to focus on understanding how to identify the domain and range of a quadratic function.

Item#	Rationale	
51	Option C is correct	To determine which system of equations can be used to find the prices in dollars of each large candle and each small candle at the store, the student could have written one equation to represent the total amount paid by a first customer for large and small candles and one equation to represent the total amount paid by a second customer for large and small candles. The total amount of 64 dollars paid by the first customer is equal to 3 times the price in dollars of each large candle, x, and 4 times the price in dollars of each small candle, y, which can be represented by the equation $3x + 4y = 64$. The total amount of 68 dollars (4 dollars more than the total amount paid by the first customer) paid by the second customer is equal to 1 time the price in dollars of each large candle, x, and 8 times the price in dollars of each small candle, y, which can be represented by the equation $x + 8y =$

Item #		Rationale
52	Option G is correct	To determine which graph best represents the function $h(x)$, the student could have first identified the linear parent function as $f(x) = x$. The student could have then realized that multiplying x by -1 means that the slope (steepness of a straight line when graphed on a coordinate grid; $m = \frac{y_2 - y_1}{x_2 - x_1}$) of the linear parent function is multiplied by -1, resulting in $-1x$, or $-x$. The student could have then realized the $+ 3$ means that the graph of linear parent function f is shifted up 3 units on the coordinate grid. The y-intercept (value where a line crosses the y-axis) of the graph of f, point $(0, 0)$, is shifted up to point $(0, 3)$, the y-intercept of the graph of h. The graph of $h(x)$ will go through the point $(0, 3)$ and have a slope of -1, which is best represented on this graph. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect	The student identified the graph of a function that has the correct y-intercept but a slope of 1 instead of -1. The student needs to focus on understanding how transformations affect the slope of the graph of a line.
	Option H is incorrect	The student identified the graph of a function that has the correct slope but a y-intercept of -3 instead of 3. The student needs to focus on understanding how transformations affect the y-intercept of the graph of a line.
	Option J is incorrect	The student identified the graph of a function that has a slope of 1 instead of -1 and a y-intercept of -3 instead of 3. The student needs to focus on understanding how transformations affect the slope and y-intercept of the graph of a line.

2019 STAAR Algebra I Rationales

Item #	Rationale	
53	Option D is correct	To determine the equivalent expression, the student could have rewritten $4\sqrt{147}$ as $4 \times \sqrt{49} \times \sqrt{3}$ and then calculated the square root (a value that, when multiplied by itself, is equal to the number under the $\sqrt{}$) of 49 to get $4 \times 7 \times \sqrt{3}$. The student could have then multiplied 4 and 7, resulting in $28\sqrt{3}$, or $28\sqrt{3}$. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely simplified $4\sqrt{147}$ as $4 \times 49 \times \sqrt{3}$, resulting in $196\sqrt{3}$, or $196\sqrt{3}$. The student needs to focus on understanding how to simplify square roots.
	Option B is incorrect	The student likely simplified O451 Tm (196)Tj /T1_4648.3n7D 38 incorrect

Item #	Rationale	
54	Option G is correct.	To determine the time, in seconds, the toy robot moves when the total distance is 408 centimeters, the student could have realized that when two variables (symbols used to represent unknown numbers), x (representing the time, in seconds) and y (representing the total distance, in centimeters), vary directly with each other, their relation can be represented by the equation $y = kx$. This equation represents the value of y, which can be determined by multiplying the value of x by the value of k (known as the constant of proportionality). The student could have calculated what number k, when multiplied by 11, equals 264, resulting in $264 = k \cdot 11 \Rightarrow \frac{264}{11} = k \Rightarrow 24 = k$. The student could have then substituted 24 for k and 408 for y in the equation $y = kx$, resulting in $408 = 24x$. To solve for x, the student could have divided both sides of the equation by 24, resulting in $17 = x$, which represents 17 seconds. The rationale for the correct answer is an efficient way to solve the problem. However, other methods could be used to solve the problem correctly.
	Option F is incorrect.	The student likely identified the value of k (known as the constant of proportionality) in the equation $y = kx$ as time, in seconds, meaning 24 seconds. The student needs to focus on understanding how to set up and solve direct variation problems.
	Option H is incorrect.	The student likely determined the time, in seconds, by calculating $\frac{408 - 4 \cdot 264}{11}$ and rounding to the nearest second, resulting in a time of 13 seconds. The student needs to focus on understanding how to set up and solve direct variation problems.
	Option J is incorrect.	The student likely determined the time, in seconds, by calculating $\frac{408}{11}$ and rounding to the nearest second, resulting in a time of 37 seconds. The student needs to focus on understanding how to set up and solve direct variation problems.