2024 STAAR Grade 7 Mathematics Rationales

Item #	Rationale
1	Option C is correct. To determine the

Item #	Rationale
4	Option B is correct To determine the volume (amount of three-dimensional space) of the triangular pyramid in cubic inches, the student should have used the formula for the volume of a pyramid, $V = \frac{1}{3}Bh$, where V is the volume, B is the area (amount of two-dimensional space covered by a surface) of the triangular base, and h is the height (vertical distance from top to bottom) of the pyramid.

2024 STAAR Grade 7 Mathematics Rationales

Item #	Rationale
5	Option D is correct. To determine the amount of milk in gallons

Item #	Rationale
6	Action, Nonfiction, Fantasy To determine which type of book is more than twice as likely to

Item #		Rationale
7	Option A is correct	To determine the value of x, the student should have set up an equation where the measures of all three angles total 180 degrees. The student should have recognized that m∠A = x = ð, m∠B = 2(x = ð • Um∠C = 2(x = ðX • The student should have then written an equation with the sum of the three angles equal to 180: x = ð = =î ðx• == î ðx180. The student should have solved this equation to find the value of x: x = ð x== î ô x== î ô A; 5xô=ì î ì =180; 5x = 160; x = 32. This is an efficient way to solve the problem; however, other methods could be used to solve the problem correctly.

Item #	Rationale
8	Option D is correct. To determine the constant of proportionality that relates y to x, the

Item #	Rationale	
9	Option A is correct	To determine the measurement closest to the height of the building in meters, the student could have set up and solved the proportion (comparison of two ratios) $\frac{x}{15} = \frac{5}{7.62}$, which compares the ratio of the height of the building in meters, x, to the height of the building in feet, 15, and the conversion factor where 1 meter is approximately equal to 3.28 feet. To solve the proportion, the student could have multiplied by each denominator (the bottom number of a fraction) on both sides of the equation, resulting in $x(3.28) = 15(1)$ or $3.28x = 15$. Finally, the student could have divided both sides of the equation by 3.28, resulting in $x \approx 4.573$. The student could have concluded that the measurement closest to the height of the building is 4.57 meters. This is an efficient way to solve the problem; however, other methods could be used to solve the problem correctly.
	Option B is incorrect	The student likely

Item #		Rationale
10	$x > -3$, $x > 9$	To determine the solution set for each inequality, the student could have used inverse operations to isolate the variable (symbol used to represent an unknown number) x on one side of the inequality. For the first inequality, $x + 3 > 12$, the student could have subtracted 9 from both sides of the inequality, yielding $x + 3 - 9 > 12 - 9$. Then the student could have simplified both sides of the inequality, yielding $x > 3$. For the second inequality, $x - 24 > 12$, the student first could have added 24 to both sides of the inequality, yielding the equivalent inequality $x > 36$. Then the student could have divided both sides of the inequality by 4, yielding $x > 9$. This is an efficient way to solve the problem; however, other methods could

Item #	Rationale	
11	Option C is correct	To determine how many more people use a car than ride a bicycle to get to work, the student could have determined how many people use a car and how many people ride a bicycle and then subtracted the number of bicycle riders from the number of car users to find the difference. To determine how many people use a car, the student could have multiplied the percentage of car users shown in the circle graph, 38%, by the number of people surveyed, 150. This results in $(38\%)(150) = (0.38)(150) = 57$, so there are 57 car users. To determine how many people ride a bicycle, the student could have determined the percentage of bicycle riders by adding up the percentages in the other categories and subtracting from 100%, resulting in a value of $100\% - (20\% + 32\% + 28\%) = 100\% - 90\% = 10\%$. Then the student could have multiplied the percent of bicycle riders by the number of people who were surveyed, 150. This results in $(10\%)(150) = (0.10)(150) = 15$, so there are 15 bicycle riders. Subtracting these values yields $57 - 15 = 42$, so the student should have concluded that there are 42 more car users than bicycle riders. This is an efficient way to solve the problem; however, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely determined the total number of bicycle riders instead of the difference between the numbers of bicycle riders and car users. To find the percentage of bicycle riders, the student likely added up the percentages from the other categories and subtracted from 100%, resulting in $100\% - (20\% + 32\% + 28\%) = 100\% - 90\% = 10\%$. Then to find the total number of bicycle riders, the student likely multiplied the percentage of bicycle riders, 10%, by the number of people in the survey, 150, resulting in $(10\%)(150) = (0.10)(150) = 15$. The student needs to focus on answering the question that is asked in a multistep scenario.
	Option B is incorrect	The student likely determined the difference between the percentages of car users and bicycle riders instead of the difference between the total numbers of car users and bicycle riders. To find the percentage of bicycle riders, the student likely added up the percentages from the other categories and subtracted from 100%, resulting in $100\% - (20\% + 32\% + 28\%) = 100\% - 90\% = 10\%$. The student then likely subtracted this value from the percentage of car users, resulting in $38\% - 10\% = 28\%$. The student needs to focus on understanding the part-to-whole relationship between percentage and sample size.
	Option D is incorrect	The student likely determined the total number of people who use a car to

Item #		Rationale
12	7,800 and any equivalent values are correct	To determine the area (amount of space covered by a surface) of the top of the table in square inches, the student could have calculated the sum of the areas of the shapes that make up the figure. The student could have divided the figure into 2 congruent trapezoids that each have a height of 30 inches and bases (lengths of the parallel sides of the trapezoid) of 130 inches and 100 inches, and 1 square with a side length of 30 inches. The student could have calculated the area of 1 trapezoid by substituting $b_1 = 130$, $b_2 = 100$, and $h = 30$ into the formula for the area of a trapezoid, $A = \frac{1}{2}(b_1 + b_2)h$, where b_1 and b_2 represent the bases of the trapezoid and h represents the height. This results in $A = \frac{1}{2}(130 + 100)(30) = 3{,}450$ square inches. Next, the student could have calculated the area of the square by using the formula $A = bh$, where b represents the base of the square and h represents the height of the square, resulting in $A = 30 \times 30 = 900$ square inches. Last, the student could have added the areas of the 2 trapezoids and the 1 square to determine the total area of the top of the table, resulting in $2(3{,}450) + 900 = 7{,}800$ square inches. This is an efficient way to solve the problem; however, other methods could be used to solve the problem correctly.

2024 STAAR Grade 7 Mathematics Rationales

Item #	Rationale

2024 STAAR Grade 7 Mathematics Rationales

Item #	Rationale

Item #	Rationale	
20	Option B is correct	To determine the final cost of groceries after the 25% off coupon was applied the student could have subtracted the discounted amount from the cost of the groceries. The student could have multiplied the cost of the groceries, $210.00, by the coupon amount, 25%, to determine the discounted amount, resulting in $(25\%)(210) = 0.25(210) = 52.50$. The student then could have subtracted this amount from the original cost, resulting in $210 - 52.50 = 157.50$. This is an efficient way to solve the problem; however, other methods could be used to solve the problem correctly.
	Option A is incorrect	The student likely determined the discounted amount from the coupon

Item #	Rationale
21	Option B is correct To determine which statement is best supported by the data in the dot plot (graph that uses dots to display data), the student could have first

Item #	Rationale
23	

Item #	Rationale
24	Option C is correct. To determine the number of days that it snowed, the student could have subtracted the total number of sunny

Item #	Rationale
25	Option D is correct To determine which measurement is closest to the area (amount of space covered by a surface) in square feet of the circular part of the lawn that is watered by the sprinkler, the student should have used the formula for the area of a circle, $A = \pi r^2$, where A is the area of the circle and r is the radius (distance from the center of the circle to a point on the circle). The student should have recognized that the radius of the circle is half the labeled diameter (length of a line segment that goes through the center of a circle and connects two points on the circle) of 14 feet, resulting in $14 \div 2 = 7$ feet. The student should then have substituted $r = 7$ into the formula for the area of a circle, resulting in $A = \pi(7^2)$, $A \approx 3.14(49)$, and $A \approx 154$ square feet. This is an efficient way to solve the problem; however, other methods could be used to solve the problem correctly.

Item #		Rationale
28	Not True; True; True Not True	To determine which equations are true when $m = -5$, the student could have substituted -5 for m to determine whether it makes a true statement. When -5 is substituted for m in the first equation, $4m - 6 = 14$, the result is $4(-5) - 6 = 14$, or $-26 = 14$. When m in $14 - \frac{\quad}{\quad} - 5)$ 2(14) Tj [14 0 Ta0

Item #	Rationale
29	Option D is correct. To determine which statement about a randomly selected student is true

Item #	Rationale	
30	Selected 2 points on the line $y = \frac{7}{6}x$, such as (0, 0) and (2, 3). Any other points on the line $y = \frac{7}{6}x$ are correct.	To determine two points on the graph that represents y, the number of cups of nuts required for x cups of dried fruit in the trail mix recipe, the student could have recognized that the rate of change (ratio of the change in y-values to the change in x-values) for this situation is $\frac{7}{6}$, since the recipe uses $1\frac{5}{6}$ cups of nuts for every cup of dried fruit. The student should have also recognized that the amount of nuts used is represented by y, and the amount of dried fruit used is represented by x. The student then could have graphed ordered pairs in which each y-value is the result of multiplying the corresponding x value by $\frac{7}{6}$, satisfying the equation $y = \frac{7}{6}x$. The graph contains the points (0, 0), (1, $1\frac{5}{6}$), (2, 3), (3, $4\frac{5}{6}$),

Item #	Rationale
31	Option C is correct. To determine the experimental probability (how likely it is that an event wi occur based on the results of an experiment) that the next card drawn is red or orange, the student should have recognized that

Item #	Rationale
33	Option C is correct. To determine which expression best represents the value of x, the student should have understood that x

Item #	Rationale
34	Option D is correct To determine the solution to the equation $4 + \frac{5}{6}T = 8 = 12$, the student could have first } u] v š Z š Œ ‰ ð } v ð š Z o (š •] } (, š Z ‹ µ resulting in $\frac{5}{6}T = 4 = 12$. Then the student could have added 4 to both sides of the equation, resulting in $\frac{5}{6}T = 4 + 4 = 12 + 4$, or $\frac{5}{6}T = 16$. Finally, the student could have multiplied both sides of the equation by 2, resulting in $2 \cdot \frac{5}{6}T = 2(16)$, or $x = 32$ This is an efficient way to solve the problem 4 0d>; hw 3.587

Item #		Rationale
35	music; physical education	To determine which inference (a conclusion based on evidence) about the preferred nonacademic subjects of sixth graders at lunch is best supported by the information in the comparative bar graph, the student should have identified the numbers of sixth graders who chose art (9), physical education (9), and music (12) and concluded that more sixth graders would choose music than would choose either one of the other two subjects. To determine which inference about the preferred nonacademic subjects of seventh graders at lunch is best supported by the information in the comparative bar graph, the student should have identified the number of seventh graders who chose art (10), physical education (14), and music (6) and concluded that more seventh graders would choose physical education than would choose either one of the other two subjects. This is an efficient

Item #	Rationale	
38	Option C is correct	To determine which inference (a conclusion based on evidence) about the number of sandwiches in each order that will be placed next Monday is best supported by the information in the dot plot (a graphical way of showing the frequency of an event by placing a dot or dots above the value on a number line), the student should have compared the total number of orders for 1 or 2 sandwiches to the total number of orders for 9 or 10 sandwiches. The student should have recognized there were 7 orders for 1 sandwich, 6 orders for 2 sandwiches, 0 orders for 9 sandwiches, and 5 orders for 10 sandwiches. The student then should have determined that the total number of orders for 1 or 2 sandwiches is ó 6 = = 13, which is more than the total number of orders for 9 or 10 sandwiches ì, = ñ A ñ X d Z Œ (} Œ U š Z • šuded v š • Z } µ o that the information in the dot plot supports the inference that more orders will be for 1 or 2 sandwiches than for 9 or 10 sandwiches.
	Option A is incorrect	The student likely multiplied the number of sandwiches in each order by the number of orders to find the total number of sandwiches ordered n BT 0qTu12 (eo10.

attending to the details of the answer options in problems that require the student to make an inf (k)11nf un